



Contents lists available at ScienceDirect

European Journal of Combinatorics

journal homepage: www.elsevier.com/locate/ejc

Some weak indivisibility results in ultrahomogeneous metric spaces

L. Nguyen Van Thé^{a,b}, N.W. Sauer^a^a Department of Mathematics and Statistics, University of Calgary, 2500 University Drive NW, Calgary, Alberta, Canada, T2N1N4^b Institut de mathématiques, Université de Neuchâtel, Emile-Argand 11, 2000 Neuchâtel, Switzerland

ARTICLE INFO

Article history:

Received 25 November 2008

Accepted 21 July 2009

Available online 18 February 2010

ABSTRACT

We study the validity of a partition property known as weak indivisibility for the integer and the rational Urysohn metric spaces. We also compare weak indivisibility to another partition property, called age-indivisibility, and provide an example of a countable ultrahomogeneous metric space which is age-indivisible but not weakly indivisible.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

The purpose of this article is the study of certain partition properties of particular metric spaces, called *ultrahomogeneous*. A metric space \mathbf{X} is ultrahomogeneous when every isometry between finite metric subspaces of \mathbf{X} can be extended to an isometry of \mathbf{X} onto itself. For example, when seen as a metric space, any Euclidean space \mathbb{R}^n has this property. So do the separable infinite dimensional Hilbert space ℓ_2 and its unit sphere \mathbb{S}^∞ . Another less known example of ultrahomogeneous metric space, though recently a well-studied object (see [5]), is the *Urysohn space*, denoted as \mathbf{U} : up to isometry, it is the unique complete separable ultrahomogeneous metric space into which every separable metric space embeds. (Here and in the rest of the paper, all the embeddings are *isometric*, that is, distance preserving.) This space also admits numerous countable analogs. For example, for various countable sets S of positive reals (see [1] for the precise condition on S), there is, up to isometry, a unique countable ultrahomogeneous metric space into which every countable metric space with distances in S embeds. When $S = \mathbb{Q}$ or \mathbb{N} this gives rise to the spaces denoted respectively as $\mathbf{U}_{\mathbb{Q}}$ (the *rational Urysohn space*) and $\mathbf{U}_{\mathbb{N}}$ (the *integer Urysohn space*). Recently, separable ultrahomogeneous metric spaces have been the center of active research because of a remarkable connection between their combinatorial behavior when submitted to finite partitions and the dynamical properties of their

E-mail addresses: nguyen@math.ucalgary.ca (L. Nguyen Van Thé), nsauer@math.ucalgary.ca (N.W. Sauer).

isometry group. For example, consider the following result. Following the terminology used in [2], we call a metric space $\mathbf{Z} = (Z, d^Z)$ *age-indivisible* if for every finite metric subspace \mathbf{Y} of \mathbf{Z} and every partition $Z = B \cup R$ of the underlying set Z of \mathbf{Z} (thought as a coloring of the points of Z with two colors, blue and red), the space \mathbf{Y} embeds in B or R . (Here and in the whole paper, boldface characters will refer to metric structures while lightface characters will refer to the corresponding underlying sets.)

Theorem (Folklore). *The spaces $\mathbf{U}_{\mathbb{Q}}$ and $\mathbf{U}_{\mathbb{N}}$ are age-indivisible.*

There are at least two directions for possible generalizations. First, one may ask what happens if instead of coloring the points of, say, the space $\mathbf{U}_{\mathbb{Q}}$, we color the isometric copies of a fixed finite metric subspace \mathbf{X} of $\mathbf{U}_{\mathbb{Q}}$. We will not touch this subject here but Kechris, Pestov and Todorcevic showed in [3] that the answer to this question (obtained by Nešetřil in [8]) has spectacular consequences on the groups $\text{iso}(\mathbf{U}_{\mathbb{Q}})$ and $\text{iso}(\mathbf{U})$ of surjective self-isometries of $\mathbf{U}_{\mathbb{Q}}$ and \mathbf{U} . For example, every continuous action of $\text{iso}(\mathbf{U})$ (equipped with the pointwise convergence topology) on a compact topological space admits a fixed point.

Another direction of generalization is to ask whether any of those spaces is *indivisible*, that is, whether B or R necessarily contains not only a copy of a fixed finite \mathbf{Y} but of the whole space itself. However, it is known that any indivisible metric space must have a bounded distance set. Therefore, the spaces $\mathbf{U}_{\mathbb{Q}}$ and $\mathbf{U}_{\mathbb{N}}$ are not indivisible. Still, in this article, we investigate whether despite this obstacle, a partition result weaker than indivisibility but stronger than age-indivisibility holds. Again, following [2], we call a metric space \mathbf{X} *weakly indivisible* when for every finite metric subspace \mathbf{Y} of \mathbf{X} and every partition $X = B \cup R$, either \mathbf{Y} embeds in B or \mathbf{X} embeds in R . Building on techniques developed in [6,11], we prove:

Theorem 1. *The space $\mathbf{U}_{\mathbb{N}}$ is weakly indivisible.*

As for $\mathbf{U}_{\mathbb{Q}}$, we are not able to prove or disprove weak indivisibility but we obtain the following weakening as a consequence of Theorem 1. If \mathbf{X} is a metric space, $Y \subset X$ and $\varepsilon > 0$, $(Y)_{\varepsilon}$ denotes the set

$$(Y)_{\varepsilon} = \{x \in X : \exists y \in Y d^{\mathbf{X}}(x, y) \leq \varepsilon\}.$$

Theorem 2. *Let $\mathbf{U}_{\mathbb{Q}} = B \cup R$ and $\varepsilon > 0$. Assume that there is a finite metric subspace \mathbf{Y} of $\mathbf{U}_{\mathbb{Q}}$ that does not embed in B . Then $\mathbf{U}_{\mathbb{Q}}$ embeds in $(R)_{\varepsilon}$.*

Another consequence of Theorem 1 is the following partition result for \mathbf{U} .

Theorem 3. *Let $\mathbf{U} = B \cup R$ and $\varepsilon > 0$. Assume that there is a compact metric subspace \mathbf{K} of \mathbf{U} that does not embed in $(B)_{\varepsilon}$. Then \mathbf{U} embeds in $(R)_{\varepsilon}$.*

Note that these results do not answer the following: for a countable ultrahomogeneous metric space is weak indivisibility a strictly stronger property than age-indivisibility? In the last section of this paper, we indicate an example of a countable ultrahomogeneous metric space which might be age-indivisible but not weakly indivisible. To our knowledge, this could be one of the first two known examples of a countable ultrahomogeneous relational structure witnessing that weak indivisibility and age-indivisibility are distinct properties (the other example will appear in [4]). Let $\mathcal{E}_{\mathbb{Q}}$ be the class of all finite metric spaces \mathbf{X} with distances in \mathbb{Q} which embed isometrically into the unit sphere S^{∞} of ℓ_2 with the property that $\{0_{\ell_2}\} \cup \mathbf{X}$ is affinely independent. It is known that there is a unique countable ultrahomogeneous metric space $S_{\mathbb{Q}}^{\infty}$ whose class of finite metric subspaces is exactly $\mathcal{E}_{\mathbb{Q}}$, and that the metric completion of $S_{\mathbb{Q}}^{\infty}$ is S^{∞} (for a proof, see [9] or [10]).

Theorem 4. *The space $S_{\mathbb{Q}}^{\infty}$ is age-indivisible.*

The proof of this result requires the use of a deep theorem due to Matoušek and Rödl in Euclidean Ramsey theory. As for the negation of weak indivisibility of $S_{\mathbb{Q}}^{\infty}$, its proof is conditioned by the validity of a strong form of the Odell–Schlumprecht distortion theorem in Banach space theory, see Section 5 for more details.

The paper is organized as follows. In Section 2, we prove [Theorem 1](#). In Section 3, we prove [Theorem 3](#). [Theorem 2](#) is proved in Section 4, and [Theorem 4](#) is proved in Section 5, where a discussion of weak indivisibility of $\mathbb{S}_{\mathbb{Q}}^{\infty}$ is also included.

2. Proof of Theorem 1

The purpose of this section is to prove [Theorem 1](#). In fact, we prove a slightly stronger result. We mentioned in introduction that there are various countable sets S of positive reals for which there is, up to isometry, a unique countable ultrahomogeneous metric space into which every countable metric space with distances in S embeds. It can be proved that when $p \in \mathbb{N}$, the integer interval $\{1, \dots, p\}$ is such a set. The corresponding countable ultrahomogeneous metric space is denoted as \mathbf{U}_p .

Theorem 5. *Let $U_{\mathbb{N}} = B \cup R$. Assume that there is $p \in \mathbb{N}$ such that \mathbf{U}_p does not embed in B . Then $\mathbf{U}_{\mathbb{N}}$ embeds in R .*

This section is devoted to the proof of [Theorem 5](#). We fix $p \in \mathbb{N}$ as well as a partition $U_{\mathbb{N}} = B \cup R$ such that \mathbf{U}_p does not embed in B . Our goal is to prove that $\mathbf{U}_{\mathbb{N}}$ embeds into R . Let $m := \lceil p/2 \rceil$ (the least integer larger than or equal to $p/2$). Recall that if $Y \subset U_{\mathbb{N}}$, the set $(Y)_{\varepsilon}$ is defined by

$$(Y)_{\varepsilon} = \{x \in U_{\mathbb{N}} : \exists y \in Y d^{\mathbf{U}_{\mathbb{N}}}(x, y) \leq \varepsilon\}.$$

In particular, if $x \in U_{\mathbb{N}}$, the set $(\{x\})_{m-1}$ denotes the set of all elements of $U_{\mathbb{N}}$ at distance $\leq m-1$ from x . We are going to construct $\tilde{U} \subset R$ isometric to $\mathbf{U}_{\mathbb{N}}$ recursively such that for every $x \in \tilde{U}$,

$$(\{x\})_{m-1} \cap \tilde{U} \subset R.$$

More precisely, fix an enumeration $\{x_n : n \in \mathbb{N}\}$ of $U_{\mathbb{N}}$. We are going to construct $\{\tilde{x}_n : n \in \mathbb{N}\} \subset U_{\mathbb{N}}$ recursively together with a decreasing sequence $\mathbf{D}_0, \mathbf{D}_1, \dots$ of metric subspaces of $\mathbf{U}_{\mathbb{N}}$ such that $x_n \mapsto \tilde{x}_n$ is an isometry and, for every $n \in \mathbb{N}$, each \mathbf{D}_n is isometric to $\mathbf{U}_{\mathbb{N}}$, $\{\tilde{x}_k : k \leq n\} \subset \mathbf{D}_n$, and $(\{\tilde{x}_n\})_{m-1} \cap \mathbf{D}_n \subset R$. To do so, we will need the notion of *Katětov map* as well as several technical lemmas.

Definition 1. Given a metric space $\mathbf{X} = (X, d^{\mathbf{X}})$, a map $f : X \rightarrow (0, +\infty)$ is *Katětov over \mathbf{X}* when

$$\forall x, y \in X, \quad |f(x) - f(y)| \leq d^{\mathbf{X}}(x, y) \leq f(x) + f(y).$$

Equivalently, one can extend the metric $d^{\mathbf{X}}$ to $X \cup \{f\}$ by defining, for every x, y in X , $\widehat{d^{\mathbf{X}}}(x, f) = f(x)$ and $\widehat{d^{\mathbf{X}}}(x, y) = d^{\mathbf{X}}(x, y)$. The corresponding metric space is then written $\mathbf{X} \cup \{f\}$. The set of all Katětov maps over \mathbf{X} is written $E(\mathbf{X})$. For a metric subspace \mathbf{X} of \mathbf{Y} and a Katětov map $f \in E(\mathbf{X})$, the *orbit of f in \mathbf{Y}* is the set $O(f, \mathbf{Y})$ defined by

$$O(f, \mathbf{Y}) = \{y \in Y : \forall x \in X d^{\mathbf{Y}}(y, x) = f(x)\}.$$

Any element $y \in O(f, \mathbf{Y})$ is said to *realize f over \mathbf{X}* . Here, the concepts of Katětov map and orbit are relevant because of the following standard reformulation of the notion of ultrahomogeneity, which will be used in the rest of the paper:

Lemma 1. *Let \mathbf{X} be a countable metric space. Then \mathbf{X} is ultrahomogeneous iff for every finite subspace \mathbf{F} of \mathbf{X} and every Katětov map f over \mathbf{F} , if $\mathbf{F} \cup \{f\}$ embeds into \mathbf{X} , then $O(f, \mathbf{X}) \neq \emptyset$.*

Proof. Postponed to Section 2.3. \square

Lemma 2. *Let G be a finite subset of $U_{\mathbb{N}}$, and g a Katětov map with domain G and with values in \mathbb{N} . Then there exists an isometric copy \mathbf{C} of $\mathbf{U}_{\mathbb{N}}$ inside $\mathbf{U}_{\mathbb{N}}$ such that:*

- (i) $G \subset \mathbf{C}$,
- (ii) $O(g, \mathbf{C}) \subset B$ or $O(g, \mathbf{C}) \subset R$.

In words, [Lemma 2](#) asserts that going to a subcopy of $\mathbf{U}_{\mathbb{N}}$ if necessary, we may assume that the orbit of g is completely included in one of the parts of the partition. Observe that as a metric space,

the orbit $O(g, \mathbf{C})$ is isometric to \mathbf{U}_n where $n = 2 \min g$ (indeed, it is countable ultrahomogeneous with distances in $\{1, \dots, n\}$ and embeds every countable metric space with distances in $\{1, \dots, n\}$).

Proof. The proof of Lemma 2 can be found in [11]. More precisely, Lemma 2 can be obtained by combining Lemmas 2 and 3 [11] after having replaced \mathbf{U}_p by $\mathbf{U}_{\mathbb{N}}$ in these statements. The proof of Lemma 3 [11] is elementary, while the proof of Lemma 2 [11] represents the core of [11] and is too lengthy to be presented here. These two proofs can be carried out with no modification once \mathbf{U}_p has been replaced by $\mathbf{U}_{\mathbb{N}}$. \square

Lemma 3. Let $G_0 \subset G$ be finite subsets of $U_{\mathbb{N}}$, and let \mathcal{G} a finite family of Katětov maps with domain G and such that for all $g, g' \in \mathcal{G}$:

$$\begin{aligned}\max(|g - g'| \upharpoonright G_0) &= \max |g - g'|, \\ \min((g + g') \upharpoonright G_0) &= \min(g + g'), \\ \min(g \upharpoonright G_0) &= \min(g).\end{aligned}$$

Then there exists an isometric copy \mathbf{C} of $\mathbf{U}_{\mathbb{N}}$ inside $\mathbf{U}_{\mathbb{N}}$ such that:

- (i) $G \cap \mathbf{C} = G_0$,
- (ii) $\forall g \in \mathcal{G} \ O(g \upharpoonright G_0, \mathbf{C}) \subset O(g, \mathbf{U}_{\mathbb{N}})$.

Proof. Postponed to Section 2.4. \square

2.1. Construction of \tilde{x}_0 and \mathbf{D}_0

First, pick an arbitrary $u \in U_{\mathbb{N}}$ and consider the map $g : \{u\} \rightarrow \mathbb{N}$ defined by $g(u) = m$. By Lemma 2, find an isometric copy \mathbf{C} of $\mathbf{U}_{\mathbb{N}}$ inside $\mathbf{U}_{\mathbb{N}}$ such that:

- (i) $u \in \mathbf{C}$,
- (ii) $O(g, \mathbf{C}) \subset B$ or $O(g, \mathbf{C}) \subset R$.

Note that since g has minimum m , the orbit $O(g, \mathbf{C})$ is isometric to \mathbf{U}_{2m} and therefore contains a copy of \mathbf{U}_p . Hence, because \mathbf{U}_p does not embed in B , the inclusion $O(g, \mathbf{C}) \subset B$ is excluded and we really have $O(g, \mathbf{C}) \subset R$. Let $\tilde{x}_0 \in O(g, \mathbf{C})$ and for every $k \leq m$ let $g_k : \{u, \tilde{x}_0\} \rightarrow \mathbb{N}$ be such that $g_k(u) = m$ and $g_k(\tilde{x}_0) = k$. The sets $G_0 = \{\tilde{x}_0\}$ and $G = \{u, \tilde{x}_0\}$, and the family $\mathcal{G} = \{g_k : k \leq m\}$ satisfy the hypotheses of Lemma 3, which allows us to obtain an isometric copy \mathbf{D}_0 of $\mathbf{U}_{\mathbb{N}}$ inside \mathbf{C} such that:

- (i) $\{u, \tilde{x}_0\} \cap \mathbf{D}_0 = \{\tilde{x}_0\}$,
- (ii) $\forall k \leq m \ O(g_k \upharpoonright \{\tilde{x}_0\}, \mathbf{D}_0) \subset O(g_k, \mathbf{C})$.

Note that for every $k \leq m$, we have $O(g_k, \mathbf{C}) \subset O(g, \mathbf{C}) \subset R$. Therefore, in \mathbf{D}_0 , all the spheres around \tilde{x}_0 with radius $k \leq m$ are included in R . So

$$(\{\tilde{x}_0\})_{m-1} \cap \mathbf{D}_0 \subset R.$$

2.2. Induction step

Assume that we constructed $\{\tilde{x}_k : k \leq n\} \subset U_{\mathbb{N}}$ together with a decreasing sequence $\mathbf{D}_0, \dots, \mathbf{D}_n$ of metric subspaces of $\mathbf{U}_{\mathbb{N}}$ such that $x_k \mapsto \tilde{x}_k$ is an isometry (recall that $\{x_n : n \in \mathbb{N}\}$ is the enumeration of $U_{\mathbb{N}}$ we fixed at the beginning of the proof), each \mathbf{D}_k is isometric to $\mathbf{U}_{\mathbb{N}}$, $\{\tilde{x}_k : k \leq n\} \subset \mathbf{D}_n$ and $(\{\tilde{x}_k\})_{m-1} \cap \mathbf{D}_n \subset R$ for every $k \leq n$. We are going to construct \tilde{x}_{n+1} and \mathbf{D}_{n+1} . Consider the map $f : \{\tilde{x}_0, \dots, \tilde{x}_n\} \rightarrow \mathbb{N}$ where

$$\forall k \leq n \ f(\tilde{x}_k) = d^{\mathbf{U}_{\mathbb{N}}}(x_k, x_{n+1}).$$

Recalling that $E(\{\tilde{x}_0, \dots, \tilde{x}_n\})$ denotes the set of all Katětov maps from the set $\{\tilde{x}_0, \dots, \tilde{x}_n\}$ to \mathbb{N} , consider the set \mathcal{G} defined by

$$\{g \in E(\{\tilde{x}_0, \dots, \tilde{x}_n\}) : \forall k \leq n (|f(\tilde{x}_k) - g(\tilde{x}_k)| \leq m - 1 \text{ and } g(\tilde{x}_k) \geq m)\}.$$

This set is finite and a repeated application of Lemma 2 allows us to construct an isometric copy \mathbf{D}_{n+1} of $\mathbf{U}_{\mathbb{N}}$ inside \mathbf{D}_n such that:

- (i) $\{\tilde{x}_0, \dots, \tilde{x}_n\} \subset D_{n+1}$,
(ii) $\forall g \in \mathcal{G}, O(g, \mathbf{D}_{n+1}) \subset B$ or R .

Note that since every $g \in \mathcal{G}$ has minimum m , the orbit $O(g, \mathbf{D}_{n+1})$ is isometric to \mathbf{U}_{2m} and therefore contains a copy of \mathbf{U}_p . Because \mathbf{U}_p does not embed in B , we consequently have

$$\forall g \in \mathcal{G}, O(g, \mathbf{D}_{n+1}) \subset R.$$

Let $\tilde{x}_{n+1} \in O(f, \mathbf{D}_{n+1})$. We claim that \tilde{x}_{n+1} and \mathbf{D}_{n+1} are as required. Note that, because $\tilde{x}_{n+1} \in O(f, \mathbf{D}_{n+1})$, we have

$$\forall k \leq n \quad d^{\mathbf{U}_N}(\tilde{x}_{n+1}, \tilde{x}_k) = f(\tilde{x}_k) = d^{\mathbf{U}_N}(x_k, x_{n+1}).$$

Therefore, $x_k \mapsto \tilde{x}_k$ is an isometry. Next we prove that $(\{\tilde{x}_{n+1}\})_{m-1} \cap D_{n+1} \subset R$. Indeed, let $y \in (\{\tilde{x}_{n+1}\})_{m-1} \cap D_{n+1}$. If $d^{\mathbf{U}_N}(\tilde{x}_k, y) \geq m$ for every $k \leq n$, then the map $d^{\mathbf{U}_N}(\cdot, y)$ is in \mathcal{G} and so $y \in O(d^{\mathbf{U}_N}(\cdot, y), \mathbf{D}_{n+1}) \subset R$. Otherwise, we have $d^{\mathbf{U}_N}(\tilde{x}_k, y) < m$ for some $k \leq n$ and

$$y \in (\{\tilde{x}_k\})_{m-1} \cap D_{n+1} \subset (\{\tilde{x}_k\})_{m-1} \cap D_n \subset R. \quad \square$$

2.3. Proof of Lemma 1

The proof is standard but we detail it here for completeness. Assume that \mathbf{X} is ultrahomogeneous. Let $\varphi : \mathbf{F} \cup \{f\} \rightarrow \mathbf{X}$ be an embedding. By ultrahomogeneity of \mathbf{X} , there is an isometry ψ of \mathbf{X} onto itself such that:

$$\forall x \in F, \quad \psi(x) = \varphi(x).$$

Then, the point $\psi^{-1}(\varphi(f))$ is in $O(f, \mathbf{X})$.

For the converse, assume that $\{x_0, \dots, x_n\}$ and $\{z_0, \dots, z_n\}$ are isometric finite subspaces of \mathbf{X} and that $\varphi : x_k \mapsto z_k$ is an isometry. We wish to extend φ to an isometry of \mathbf{X} onto itself. We do that thanks to a back and forth method. First, extend $\{x_0, \dots, x_n\}$ and $\{z_0, \dots, z_n\}$ so that $\{x_k : k \in \mathbb{N}\} = \{z_k : k \in \mathbb{N}\} = X$. For $k \leq n$, let $\sigma(k) = \tau(k) = k$. Then, set $\sigma(n+1) = n+1$. Consider the map f_{n+1} defined on $\{\varphi(x_{\sigma(k)}) : k \leq n\}$ by:

$$\forall k \leq n, \quad f_{n+1}(\varphi(x_{\sigma(k)})) = d^{\mathbf{X}}(x_{\sigma(n+1)}, x_{\sigma(k)}).$$

Observe that f_{n+1} is Katětov over $\{\varphi(x_{\sigma(k)}) : k \leq n\}$ and that the space $\{\varphi(x_{\sigma(k)}) : k \leq n\} \cup \{f_{n+1}\}$ is isometric to $\{x_{\sigma(k)} : k \leq n+1\}$. By hypothesis on \mathbf{X} , we can consequently find a point in $O(f_{n+1}, \mathbf{X})$, call it $\varphi(x_{\sigma(n+1)})$. Next, set:

$$\tau(n+1) = \min\{k \in \mathbb{N} : z_k \notin \{\varphi(x_{\sigma(i)}) : i \leq n\}\}.$$

Consider the map g_{n+1} defined on $\{x_{\sigma(k)} : k \leq n\}$ by:

$$\forall k \leq n, \quad g_{n+1}(x_{\sigma(k)}) = d^{\mathbf{X}}(z_{\tau(n+1)}, \varphi(x_{\sigma(k)})).$$

Then g_{n+1} is Katětov over the space $\{x_{\sigma(k)} : k \leq n\}$ and the corresponding union $\{x_{\sigma(k)} : k \leq n\} \cup \{g_{n+1}\}$ is isometric to $\{\varphi(x_{\sigma(k)}) : k \leq n\} \cup \{z_{\tau(n+1)}\}$. So again, by hypothesis on \mathbf{X} , we can find a point in $O(g_{n+1}, \mathbf{X})$, call it $\varphi^{-1}(z_{\tau(n+1)})$. In general, if σ and τ have been defined up to m and φ has been defined on $T_m := \{x_{\sigma(0)}, \dots, x_{\sigma(m)}\} \cup \{\varphi^{-1}(z_{\sigma(0)}), \dots, \varphi(z_{\sigma(m)})\}$, set:

$$\sigma(m+1) = \min\{k \in \mathbb{N} : x_k \notin T_m\}.$$

Consider the map f_{m+1} defined on $\varphi(T_m)$ by:

$$\forall k \leq m, \quad \begin{cases} f_{m+1}(\varphi(x_{\sigma(k)})) = d^{\mathbf{X}}(x_{\sigma(m+1)}, x_{\sigma(k)}) \\ f_{m+1}(z_{\tau(k)}) = d^{\mathbf{X}}(x_{\sigma(m+1)}, \varphi^{-1}(z_{\tau(k)})). \end{cases}$$

Observe that f_{m+1} is Katětov over $\varphi(T_m)$ and that $\varphi(T_m \cup \{f_{m+1}\})$ is isometric to $T_m \cup \{x_{\sigma(m+1)}\}$. By hypothesis on \mathbf{X} , we can consequently find a point in $O(f_{m+1}, \mathbf{X})$, call it $\varphi(x_{\sigma(m+1)})$. Next, let:

$$\tau(m+1) = \min\{k \in \mathbb{N} : z_k \notin \{\varphi(x_{\sigma(i)}) : i < n+1\}\}.$$

Consider the map g_{m+1} defined on T_m by:

$$\forall k \leq m, \quad \begin{cases} g_{m+1}(x_{\sigma(k)}) = d^{\mathbf{X}}(z_{\tau(m+1)}, \varphi(x_{\sigma(k)})) \\ g_{m+1}(\varphi^{-1}(z_{\tau(k)})) = d^{\mathbf{X}}(z_{\tau(m+1)}, z_{\tau(k)}). \end{cases}$$

Then g_{m+1} is Katětov over T_m and the union $T_m \cup \{g_{m+1}\}$ is isometric to $\varphi(T_m \cup \{z_{\tau(m+1)}\})$. So again, by hypothesis on \mathbf{X} , we can find a point in $O(g_{m+1}, \mathbf{X})$, call it $\varphi^{-1}(z_{\tau(m+1)})$. After infinitely many steps, we are left with an isometry φ with domain $\mathbf{X} = \{x_k : k \in \mathbb{N}\}$ and range $\mathbf{X} = \{z_k : k \in \mathbb{N}\}$. This finishes the proof.

2.4. Proof of Lemma 3

Lemma 3 is a modified version of a result proved in [11], namely **Lemma 5**. Let $G_0 \subset G$ be finite subsets of $\mathbf{U}_{\mathbb{N}}$, \mathcal{G} a family of Katětov maps with domain G and such that for every $g, g' \in \mathcal{G}$:

$$\begin{aligned} \max(|g - g'| \upharpoonright G_0) &= \max|g - g'|, \\ \min((g + g') \upharpoonright G_0) &= \min(g + g'). \end{aligned}$$

We need to produce an isometric copy \mathbf{C} of $\mathbf{U}_{\mathbb{N}}$ inside $\mathbf{U}_{\mathbb{N}}$ such that:

- (i) $G \cap \mathbf{C} = G_0$.
- (ii) $\forall g \in \mathcal{G} \ O(g \upharpoonright G_0, \mathbf{C}) \subset O(g, \mathbf{U}_{\mathbb{N}})$.

First, observe that it suffices to provide the proof assuming that G is of the form $G_0 \cup \{z\}$. The general case is then handled by repeating the procedure.

Lemma 4. Let \mathbf{X} be a finite subspace of $\bigcup\{O(g \upharpoonright G_0, \mathbf{U}_{\mathbb{N}}) : g \in \mathcal{G}\}$. Then there is an isometry φ on $\mathbf{U}_{\mathbb{N}}$ fixing $G_0 \cup (X \cap \bigcup\{O(g, \mathbf{U}_{\mathbb{N}}) : g \in \mathcal{G}\})$ and such that:

$$\forall g \in \mathcal{G} \quad \varphi(X \cap O(g \upharpoonright G_0, \mathbf{U}_{\mathbb{N}})) \subset O(g, \mathbf{U}_{\mathbb{N}}).$$

Proof. For $x \in X$, there is a unique element $g_x \in \mathcal{G}$ such that $x \in O(g_x \upharpoonright G_0, \mathbf{U}_{\mathbb{N}})$. Let k be the map defined on $G_0 \cup X$ by

$$k(x) = \begin{cases} d^{\mathbf{U}_{\mathbb{N}}}(x, z) & \text{if } x \in G_0, \\ g_x(z) & \text{if } x \in X. \end{cases}$$

Claim 1. The map k is Katětov.

Proof. The metric space $G_0 \cup \{z\}$ witnesses that k is Katětov over G_0 . Hence, it suffices to check that for every $x \in X$ and $y \in G_0 \cup X$,

$$|k(x) - k(y)| \leq d^{\mathbf{U}_{\mathbb{N}}}(x, y) \leq k(x) + k(y).$$

Consider first the case $y \in G_0$. Then $d^{\mathbf{U}_{\mathbb{N}}}(x, y) = g_x(y)$ and we need to check that

$$|g_x(z) - d^{\mathbf{U}_{\mathbb{N}}}(y, z)| \leq g_x(y) \leq g_x(z) + d^{\mathbf{U}_{\mathbb{N}}}(y, z).$$

Or equivalently,

$$|g_x(z) - g_x(y)| \leq d^{\mathbf{U}_{\mathbb{N}}}(y, z) \leq g_x(z) + g_x(y).$$

But this is true since g_x is Katětov over $G_0 \cup \{z\}$. Consider now the case $y \in X$. Then $k(y) = g_y(z)$ and we need to check

$$|g_x(z) - g_y(z)| \leq d^{\mathbf{U}_{\mathbb{N}}}(x, y) \leq g_x(z) + g_y(z).$$

But since \mathbf{X} is a subspace of $\bigcup\{O(g \upharpoonright G_0, \mathbf{U}_{\mathbb{N}}) : g \in \mathcal{G}\}$, we have, for every $u \in G_0$,

$$|d^{\mathbf{U}_{\mathbb{N}}}(x, u) - d^{\mathbf{U}_{\mathbb{N}}}(u, y)| \leq d^{\mathbf{U}_{\mathbb{N}}}(x, y) \leq d^{\mathbf{U}_{\mathbb{N}}}(x, u) + d^{\mathbf{U}_{\mathbb{N}}}(u, y).$$

Since $x \in O(g_x \upharpoonright G_0, \mathbf{U}_{\mathbb{N}})$ and $y \in O(g_y \upharpoonright G_0, \mathbf{U}_{\mathbb{N}})$, this is equivalent to

$$|g_x(u) - g_y(u)| \leq d^{\mathbf{U}_{\mathbb{N}}}(x, y) \leq g_x(u) + g_y(u).$$

Therefore,

$$\max(|g_x - g_y| \upharpoonright G_0) \leq d^{\mathbf{U}_N}(x, y) \leq \min((g_x + g_y) \upharpoonright G_0).$$

Now, by hypothesis on \mathcal{G} , this latter inequality remains valid if G_0 is replaced by $G_0 \cup \{z\}$. The required inequality follows. \square

By ultrahomogeneity of \mathbf{U}_N (or, more precisely, by its equivalent reformulation provided in Lemma 1), we can pick a point $z' \in O(k, \mathbf{U}_N)$. The metric space $G_0 \cup (X \cap \bigcup\{O(g, \mathbf{U}_N) : g \in \mathcal{G}\}) \cup \{k\}$ being isometric to the subspace of \mathbf{U}_N supported by $G_0 \cup (X \cap \bigcup\{O(g, \mathbf{U}_N) : g \in \mathcal{G}\}) \cup \{z\}$, so is the subspace of \mathbf{U}_N supported by $G_0 \cup (X \cap \bigcup\{O(g, \mathbf{U}_N) : g \in \mathcal{G}\}) \cup \{z'\}$. By ultrahomogeneity again, we can find a surjective isometry φ of \mathbf{U}_N fixing $G_0 \cup (X \cap \bigcup\{O(g, \mathbf{U}_N) : g \in \mathcal{G}\})$ and such that $\varphi(z') = z$. Then φ is as required: let $g \in \mathcal{G}$ and $x \in O(g \upharpoonright G_0, \mathbf{U}_N)$. Then:

$$d^{\mathbf{U}_N}(\varphi(x), z) = d^{\mathbf{U}_N}(\varphi(x), \varphi(z')) = d^{\mathbf{U}_N}(x, z') = k(x) = g(z).$$

That is, $\varphi(x) \in O(g, \mathbf{U}_N)$. \square

Lemma 5. *There is an isometric embedding ψ of $G_0 \cup \bigcup\{O(g \upharpoonright G_0, \mathbf{U}_N) : g \in \mathcal{G}\}$ into $G_0 \cup \bigcup\{O(g, \mathbf{U}_N) : g \in \mathcal{G}\}$ fixing G_0 such that:*

$$\forall g \in \mathcal{G} \quad \psi(O(g \upharpoonright G_0, \mathbf{U}_N)) \subset O(g, \mathbf{U}_N).$$

Proof. Let $\{x_n : n \in \mathbb{N}\}$ enumerate $\bigcup\{O(g \upharpoonright G_0, \mathbf{U}_N) : g \in \mathcal{G}\}$. For $n \in \mathbb{N}$, let g_n be the only $g \in \mathcal{G}$ such that $x_n \in O(g_n \upharpoonright G_0, \mathbf{U}_N)$. Apply Lemma 4 inductively to construct a sequence $(\psi_n)_{n \in \mathbb{N}}$ of surjective isometries of \mathbf{U}_N such that for every $n \in \mathbb{N}$, ψ_n fixes $G_0 \cup \psi_{n-1}(\{x_k : k < n\})$ and $\psi_n(x_n) \in O(g_n, \mathbf{U}_N)$. Then ψ defined on $G_0 \cup \{x_n : n \in \mathbb{N}\}$ by $\psi \upharpoonright G_0 = id_{G_0}$ and $\psi(x_n) = \psi_n(x_n)$ is as required. \square

We now turn to the proof of Lemma 3. Let \mathbf{Y} and \mathbf{Z} be the metric subspaces of \mathbf{U}_N supported by $G \cup \bigcup\{O(g, \mathbf{U}_N) : g \in \mathcal{G}\}$ and $G_0 \cup \bigcup\{O(g \upharpoonright G_0, \mathbf{U}_N) : g \in \mathcal{G}\}$ respectively. Let $i_0 : \mathbf{Z} \rightarrow \mathbf{U}_N$ be the isometric embedding provided by the identity. By Lemma 5, the space \mathbf{Z} embeds isometrically into \mathbf{Y} via an isometry j_0 that fixes G_0 . We can therefore consider the metric space \mathbf{W} obtained by gluing \mathbf{U}_N and \mathbf{Y} via an identification of $\mathbf{Z} \subset \mathbf{U}_N$ and $j_0(\mathbf{Z}) \subset \mathbf{Y}$. The space \mathbf{W} is described in Fig. 1.

Formally, the space \mathbf{W} can be constructed thanks to a property of countable metric spaces with distances in \mathbb{N} known as *strong amalgamation*: we can find a countable metric space \mathbf{W} with distances in \mathbb{N} and isometric embeddings $i_1 : \mathbf{U}_N \rightarrow \mathbf{W}$ and $j_1 : \mathbf{Y} \rightarrow \mathbf{W}$ such that:

- $i_1 \circ i_0 = j_1 \circ j_0$,
- $W = i_1(U_N) \cup j_1(Y)$,
- $i_1(U_N) \cap j_1(Y) = (i_1 \circ i_0)(Z) = (j_1 \circ j_0)(Z)$,
- for every $x \in U_N$ and $y \in Y$:

$$\begin{aligned} d^{\mathbf{W}}(i_1(x), j_1(y)) &= \min\{d^{\mathbf{W}}(i_1(x), i_1 \circ i_0(z)) + d^{\mathbf{W}}(j_1 \circ j_0(z), j_1(y)) : z \in Z\} \\ &= \min\{d^{\mathbf{U}_N}(x, i_0(z)) + d^{\mathbf{Y}}(j_0(z), y) : z \in Z\} \\ &= \min\{d^{\mathbf{U}_N}(x, z) + d^{\mathbf{Y}}(j_0(z), y) : z \in Z\}. \end{aligned}$$

The crucial point here is that in \mathbf{W} , every $x \in i_1(U_N)$ realizing some $g \upharpoonright G_0$ over $i_1(G_0)$ also realizes g over $j_1(G)$.

Using \mathbf{W} , we show how \mathbf{C} can be constructed inductively: consider an enumeration $\{x_n : n \in \mathbb{N}\}$ of $i_1(U_N)$ admitting $i_1(G_0)$ as an initial segment. Assume that the points $\varphi(x_0), \dots, \varphi(x_n)$ are constructed so that:

- the map φ is an isometry,
- $\text{dom } \varphi \subset i_1(U_N)$,
- $\varphi(x_0), \dots, \varphi(x_n) \in U_N$,
- $\varphi(i_1(x)) = x$ whenever $x \in G_0$,
- $d^{\mathbf{U}_N}(\varphi(x_k), z) = d^{\mathbf{W}}(x_k, j_1(z))$ whenever $z \in G$ and $k \leq n$.




Fig. 1. The space \mathbf{W} .

We want to construct $\varphi(x_{n+1})$. Consider e defined on $\{\varphi(x_k) : k \leq n\} \cup G$ by:

$$\begin{cases} \forall k \leq n & e(\varphi(x_k)) = d^{\mathbf{W}}(x_k, x_{n+1}), \\ \forall z \in G & e(z) = d^{\mathbf{W}}(j_1(z), x_{n+1}). \end{cases}$$

Observe that the metric subspace of \mathbf{W} given by $\{x_k : k \leq n+1\} \cup j_1(G)$ witnesses that e is Katětov. It follows that the set $O(e, \mathbf{U}_{\mathbb{N}})$ is not empty and $\varphi(x_{n+1})$ can be chosen in it. \square

3. Proof of Theorem 3

The purpose of this section is to prove Theorem 3. So let $U = B \cup R$ and $\varepsilon > 0$. Assume that there is a compact metric subspace \mathbf{K} of \mathbf{U} that does not embed in $(B)_{\varepsilon}$. We wish to show that \mathbf{U} embeds in $(R)_{\varepsilon}$.

3.1. Proof of Theorem 3

We will use the result of Theorem 1 as well as two technical lemmas, whose proofs are postponed to Sections 3.2 and 3.3.

Lemma 6. Let $q \in \mathbb{N}$ be positive. Then there is an isometric copy $\mathbf{U}_{\mathbb{N}/q}^*$ of $\mathbf{U}_{\mathbb{N}/q}$ in \mathbf{U} such that for every subspace $\tilde{\mathbf{V}}$ of $\mathbf{U}_{\mathbb{N}/q}^*$ isometric to $\mathbf{U}_{\mathbb{N}/q}$, the set $(\tilde{\mathbf{V}})_{1/q}$ includes an isometric copy of \mathbf{U} .

The second lemma states that in \mathbf{U} , the copies of the compact space \mathbf{K} can be captured by a single finite metric subspace of \mathbf{U} :

Lemma 7. There is a finite metric subspace \mathbf{Y} of \mathbf{U} with rational distances such that \mathbf{K} embeds in $(\tilde{\mathbf{Y}})_{\varepsilon/2}$ for every subspace $\tilde{\mathbf{Y}}$ of \mathbf{U} isometric to \mathbf{Y} .

Assuming Lemmas 6 and 7, the proof of Theorem 3 goes as follows: choose $q \in \mathbb{N}$ large enough so that $1/q \leq \varepsilon$ and all distances appearing in \mathbf{Y} are integer multiples of $1/q$. The partition $U = B \cup R$ induces a partition of $\mathbf{U}_{\mathbb{N}/q}^*$ provided by Lemma 6. Note that \mathbf{Y} does not embed in B : indeed, if a subspace

$\tilde{\mathbf{Y}}$ of B were isometric to \mathbf{Y} , then $(\tilde{\mathbf{Y}})_\varepsilon \subset (B)_\varepsilon$ and by Lemma 7, the space \mathbf{K} would embed in $(B)_\varepsilon$, which is not the case. Observe now that by weak indivisibility of the space $\mathbf{U}_\mathbb{N}$ (Theorem 1), the space $\mathbf{U}_{\mathbb{N}/q}$ is weakly indivisible as well, so there is a subspace $\tilde{\mathbf{V}}$ of $\mathbf{U}_{\mathbb{N}/q}^*$ isometric to $\mathbf{U}_{\mathbb{N}/q}$ such that $\tilde{\mathbf{V}} \subset R$. By construction of $\mathbf{U}_{\mathbb{N}/q}^*$, the set $(\tilde{\mathbf{V}})_{1/q}$ includes an isometric copy $\tilde{\mathbf{U}}$ of \mathbf{U} . To complete the proof, notice that $\tilde{\mathbf{U}} \subset (\tilde{\mathbf{V}})_{1/q} \subset (\tilde{\mathbf{V}})_\varepsilon \subset (R)_\varepsilon$.

3.2. Proof Lemma 6

Lemma 6 is a modified version of a result proved in [6], whose statement appears at the very beginning of Proposition 5. Its proof is an easy modification of Lemma 2 [6] and is included here for completeness. The core of the proof is contained in Lemma 8 which we present now. Fix an enumeration $\{y_n : n \in \mathbb{N}\}$ of U_Q . For a number α , let $\lceil \alpha \rceil_q$ denote the smallest number $\geq \alpha$ of the form l/q with l integer. The function $\lceil \cdot \rceil_q$ is subadditive and nondecreasing. Hence, the composition $\lceil d^Z \rceil_q = \lceil \cdot \rceil_q \circ d^{U_Q}$ is a metric on U_Q . Let \mathbf{X}_q be the metric space $(U_Q, \lceil d^{U_Q} \rceil_q)$. The underlying set of \mathbf{X}_q is really $\{y_n : n \in \mathbb{N}\}$ but to avoid confusion, we write it $\{x_n : n \in \mathbb{N}\}$, being understood that for every $n \in \mathbb{N}$, $x_n = y_n$. On the other hand, note that $\mathbf{U}_{\mathbb{N}/q}$ and \mathbf{X}_q embed isometrically into each other: \mathbf{X}_q embeds in $\mathbf{U}_{\mathbb{N}/q}$ because any countable metric space with distances in \mathbb{N}/q embeds in $\mathbf{U}_{\mathbb{N}/q}$, and $\mathbf{U}_{\mathbb{N}/q}$ embeds in \mathbf{X}_q because any copy of $\mathbf{U}_{\mathbb{N}/q}$ in U_Q remains isometric to $\mathbf{U}_{\mathbb{N}/q}$ in $\mathbf{X}_q = (U_Q, \lceil d^{U_Q} \rceil_q)$.

Lemma 8. *There is a countable metric space \mathbf{Z} including \mathbf{X}_q such that for every strictly increasing $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ such that $x_n \mapsto x_{\sigma(n)}$ is an isometry, the set $(\{x_{\sigma(n)} : n \in \mathbb{N}\})_{1/q}$ includes an isometric copy of U_Q .*

Assuming Lemma 8, we now show how we can construct $\mathbf{U}_{\mathbb{N}/q}^*$. The space \mathbf{Z} is countable so we may assume that it is a subspace of \mathbf{U} . Now, take $\mathbf{U}_{\mathbb{N}/q}^*$ a subspace of \mathbf{X}_q and isometric to $\mathbf{U}_{\mathbb{N}/q}$. We claim that $\mathbf{U}_{\mathbb{N}/q}^*$ works: let $\tilde{\mathbf{V}}$ be a subspace of $\mathbf{U}_{\mathbb{N}/q}^*$ isometric to $\mathbf{U}_{\mathbb{N}/q}$. We first show that $(\tilde{\mathbf{V}})_{1/q}$ includes a copy of U_Q . The enumeration $\{x_n : n \in \mathbb{N}\}$ induces a linear ordering $<$ of $\tilde{\mathbf{V}}$. According to Lemma 8, it suffices to show that $(\tilde{\mathbf{V}}, <)$ includes a copy of $\{x_n : n \in \mathbb{N}\}_<$ seen as an ordered metric space. To do that, observe that since \mathbf{X}_q embeds isometrically into $\mathbf{U}_{\mathbb{N}/q}$, there is a linear ordering $<^*$ of $\mathbf{U}_{\mathbb{N}/q}$ such that $\{x_n : n \in \mathbb{N}\}_<$ embeds into $(\mathbf{U}_{\mathbb{N}/q}, <^*)$ as ordered metric space. Therefore, it is enough to show:

Claim 2. *$(\tilde{\mathbf{V}}, <)$ includes a copy of $(\mathbf{U}_{\mathbb{N}/q}, <^*)$.*

Proof. Write

$$(U_{\mathbb{N}/q}, <^*) = \{s_n : n \in \mathbb{N}\}_{<^*}$$

$$(\tilde{\mathbf{V}}, <) = \{t_n : n \in \mathbb{N}\}_{<}$$

Let $\sigma(0) = 0$. If $\sigma(0) < \dots < \sigma(n)$ are chosen such that $s_k \mapsto t_{\sigma(k)}$ is a finite isometry, observe that the following set is infinite

$$\{i \in \mathbb{N} : \forall k \leq n \ d^{U_{\mathbb{N}/q}}(t_{\sigma(k)}, t_i) = d^{U_{\mathbb{N}/q}}(s_k, s_{n+1})\}.$$

Therefore, simply take $\sigma(n+1)$ in that set and larger than $\sigma(n)$. \square

Observe that since the metric completion of U_Q is \mathbf{U} , the closure of $(\tilde{\mathbf{V}})_{1/q}$ in \mathbf{U} includes a copy of \mathbf{U} . Hence we are done since $(\tilde{\mathbf{V}})_{1/q}$ is closed in \mathbf{U} .

We now turn to the proof of Lemma 8. The strategy is first to provide the set Z where the required metric space \mathbf{Z} is supposed to be based on, and then to argue that the distance d^Z can be obtained (Lemmas 9–13). To construct Z , proceed as follows: for $t \subset \mathbb{N}$, write t as the strictly increasing enumeration of its elements:

$$t = \{t_i : i \in |t|_{<}\}.$$

Now, let T be the set of all finite nonempty subsets t of \mathbb{N} such that $x_n \mapsto x_{t_n}$ is an isometry between $\{x_n : n \in |t|\}$ and $\{x_{t_n} : n \in |t|\}$. This set T is a tree (in the order-theoretic sense) when ordered by

end-extension. Let

$$Z = X_q \dot{\cup} T.$$

For $z \in Z$, define

$$\pi(z) = \begin{cases} z & \text{if } z \in X_q, \\ x_{\max z} & \text{if } z \in T. \end{cases}$$

Now, consider an edge-labelled graph structure on Z by defining δ with domain $\text{dom}(\delta) \subset Z \times Z$ as follows:

- If $s, t \in T$, then $(s, t) \in \text{dom}(\delta)$ iff s and t are $<_T$ -comparable. In this case,

$$\delta(s, t) = d^{\mathbf{U}_Q}(y_{|s|-1}, y_{|t|-1}).$$

- If $x, y \in X_q$, then (x, y) is always in $\text{dom}(\delta)$ and

$$\delta(x, y) = d^{\mathbf{X}_q}(x, y).$$

- If $t \in T$ and $x \in X_q$, then (x, s) and (s, x) are in $\text{dom}(\delta)$ iff $x = \pi(t)$. In this case

$$\delta(x, s) = \delta(s, x) = \frac{1}{q}.$$

For a branch b of T and $i \in \mathbb{N}$, let $b(i)$ be the unique element of b with height i in T . Observe that $b(i)$ is a $(i+1)$ -element subset of \mathbb{N} . Observe also that for every $i, j \in \mathbb{N}$, $b(i)$ is connected to $\pi(b(i))$ and $b(j)$, and

- (i) $\delta(b(i), \pi(b(i))) = 1/q$,
- (ii) $\delta(b(i), b(j)) = d^{\mathbf{U}_Q}(y_i, y_j)$,
- (iii) $\delta(\pi(b(i)), \pi(b(j)))$ is equal to any of the following quantities:

$$d^{\mathbf{X}_q}(x_{\max b(i)}, x_{\max b(j)}) = d^{\mathbf{X}_q}(x_i, x_j) = \lceil d^{\mathbf{U}_Q}(y_i, y_j) \rceil_q.$$

In particular, if b is a branch of T , then δ induces a metric on b and the map from \mathbf{U}_Q to b mapping y_i to $b(i)$ is a surjective isometry. We claim that if we can show that δ can be extended to a metric d^Z on Z , then [Lemma 8](#) will be proved. Indeed, let

$$\tilde{X}_q = \{x_{\sigma(n)} : n \in \mathbb{N}\} \subset X_q,$$

with $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ strictly increasing and $x_n \mapsto x_{\sigma(n)}$ distance preserving. See $\text{ran}(\sigma)$, the range of σ , as a branch b of T . Then $(b, d^Z) = (b, \delta)$ is isometric to \mathbf{U}_Q and

$$b \subset (\pi(b))_{1/q} = (\tilde{X}_q)_{1/q}.$$

Our goal now is consequently to show that δ can be extended to a metric on Z with values in $[0, 1]$. For $x, y \in Z$, and $n \in \mathbb{N}$ strictly positive, define a *path from x to y of size n* as a finite sequence $\gamma = (z_i)_{i < n}$ such that $z_0 = x, z_{n-1} = y$ and for every $i < n-1$,

$$(z_i, z_{i+1}) \in \text{dom}(\delta).$$

For x, y in Z , let $P(x, y)$ be the set of all paths from x to y . If $\gamma = (z_i)_{i < n}$ is in $P(x, y)$, $\|\gamma\|$ is defined as:

$$\|\gamma\| = \sum_{i=0}^{n-1} \delta(z_i, z_{i+1}).$$

We are going to see that the required metric can be obtained with d^Z defined by

$$d^Z(x, y) = \inf\{\|\gamma\| : \gamma \in P(x, y)\}.$$

Equivalently, we are going to show that for every $(x, y) \in \text{dom}(\delta)$, every path γ from x to y is *metric*, i.e. satisfies the following inequality:

$$\delta(x, y) \leq \|\gamma\|. \tag{1}$$




Fig. 2. Irreducible cycles.

Let $x, y \in Z$. Call a path γ from x to y *trivial* when $\gamma = (x, y)$ and *irreducible* when no proper subsequence of γ is a non-trivial path from x to y . Finally, say that γ is a *cycle* when $(x, y) \in \text{dom}(\delta)$. It should be clear that to prove that d^Z works, it is enough to show that the previous inequality (1) is true for every irreducible cycle. Note that even though δ takes only rational values, it might not be the case for d^Z . We now turn to the study of the irreducible cycles in Z .

Lemma 9. Let $x, y \in T$. Assume that x and y are not $<_T$ -comparable. Let γ be an irreducible path from x to y in T . Then there is $z \in T$ such that $z <_T x, z <_T y$ and $\gamma = (x, z, y)$.

Proof. Write $\gamma = (z_i)_{i \leq n+1}$. z_1 is connected to x so z_1 is $<_T$ -comparable with x . We claim that $z_1 <_T x$: otherwise, $x <_T z_1$ and every element of T which is $<_T$ -comparable with z_1 is also $<_T$ -comparable with x . In particular, z_2 is $<_T$ -comparable with x , a contradiction since z_2 and x are not connected. We now claim that $z_1 <_T y$. Indeed, observe that $z_1 <_T z_2$: otherwise, $z_2 <_T z_1 <_T x$ so $z_2 <_T x$ contradicting irreducibility. Now, every element of T which is $<_T$ -comparable with z_2 is also $<_T$ -comparable with z_1 , so no further element can be added to the path. Hence $z_2 = y$ and we can take $z_1 = z$. \square

Lemma 10. Every non-trivial irreducible cycle in X_q has size 3.

Proof. Obvious since δ induces the metric d^{X_q} on X_q . \square

Lemma 11. Every non-trivial irreducible cycle in T has size 3 and is included in a branch.

Proof. Let $c = (z_i)_{i \leq n}$ be a non-trivial irreducible cycle in T . We may assume that $z_0 <_T z_{n-1}$. Now, observe that every element of T which is comparable with z_0 is also comparable with z_{n-1} . In particular, z_1 is such an element. It follows that $n = 3$ and that z_0, z_1, z_2 are in a same branch. \square

Lemma 12. Every irreducible cycle in Z intersecting both X_q and T is supported by a set whose form is one of the following ones (see Fig. 2).

Proof. Let C be a set supporting an irreducible cycle c intersecting both X_q and T . It should be clear that $|C \cap X_q| \leq 2$: otherwise since any two points in X_q are connected, c would admit a strict subcycle, contradicting irreducibility.

If $C \cap X_q$ has size 1, let z_0 be its unique element. In c , z_0 is connected to two elements which we denote z_1 and z_3 . Note that $z_1, z_3 \in T$ so $\pi(z_1) = \pi(z_3) = z_0$. Since elements in T which are connected never project on a same point, it follows that z_1, z_3 are $<_T$ -incomparable. Now, c induces an irreducible path from z_1 to z_3 in T so from Lemma 9, there is $z_2 \in C$ such that $z_2 <_T z_1, z_2 <_T z_3$, and we are in Case 2.

Assume now that $C \cap X_q = \{z_0, z_4\}$. Then there are $z_1, z_3 \in C \cap T$ such that $\pi(z_1) = z_0$ and $\pi(z_3) = z_4$. Note that since $z_0 \neq z_4$, we must have $z_1 \neq z_3$. Now, $C \cap T$ induces an irreducible path from z_1 to z_3 in T . By Lemma 9, either z_1 and z_3 are compatible and in this case, we are in Case 1, or z_1 and z_3 are $<_T$ -incomparable and there is $z_2 \in C \cap T$ such that $z_2 <_T z_1, z_2 <_T z_3$ and we are in Case 3. \square

Lemma 13. Every non-trivial irreducible cycle in Z is metric.

Proof. Let c be an irreducible cycle in Z . If c is supported by X_q , then, by Lemma 10, c has size 3 and is metric since δ induces a metric on X_q . If c is supported by T , then, by Lemma 11, c also has size 3 and is included in a branch b of T . Since δ induces a metric on b , c is metric. We consequently assume that c intersects both X_q and T . According to Lemma 12, c is supported by a set C whose form is covered by one of the Cases 1, 2 or 3. So to prove the present lemma, it is enough to show every cycle obtained from a re-indexing of the cycles described in those cases is metric.

Case 1: The required inequalities are obvious after having observed that

$$\delta(z_0, z_3) = \lceil \delta(z_1, z_2) \rceil_q \quad \text{and} \quad \delta(z_0, z_1) = \delta(z_2, z_3) = \frac{1}{q}.$$

Case 2: Notice that $\delta(z_0, z_1) = \delta(z_0, z_3) = 1/q$. So the inequalities we need to prove are

$$\delta(z_1, z_2) \leq \delta(z_2, z_3) + \frac{2}{q}, \quad (2)$$

$$\delta(z_2, z_3) \leq \delta(z_1, z_2) + \frac{2}{q}. \quad (3)$$

By symmetry, it suffices to verify that (2) holds. Observe that since $\pi(z_1) = \pi(z_3) = z_0$, we must have $\lceil \delta(z_1, z_2) \rceil_q = \lceil \delta(z_2, z_3) \rceil_q$. So:

$$\delta(z_1, z_2) \leq \lceil \delta(z_1, z_2) \rceil_q = \lceil \delta(z_2, z_3) \rceil_q \leq \delta(z_2, z_3) + \frac{2}{q}.$$

Case 3: Observe that $\delta(z_0, z_1) = \delta(z_3, z_4) = 1/q$, so the inequalities we need to prove are

$$\delta(z_1, z_2) \leq \delta(z_2, z_3) + \delta(z_0, z_4) + \frac{2}{q}, \quad (4)$$

$$\delta(z_0, z_4) \leq \delta(z_1, z_2) + \delta(z_2, z_3) + \frac{2}{q}. \quad (5)$$

For (4):

$$\begin{aligned} \delta(z_1, z_2) &\leq \lceil \delta(z_1, z_2) \rceil_q \\ &= \delta(\pi(z_1), \pi(z_2)) \\ &= \delta(z_0, \pi(z_2)) \\ &\leq \delta(z_0, z_4) + \delta(z_4, \pi(z_2)) \\ &= \delta(z_0, z_4) + \lceil \delta(z_3, z_2) \rceil_q \\ &\leq \delta(z_0, z_4) + \delta(z_2, z_3) + \frac{2}{q}. \end{aligned}$$

For (5): Write $z_1 = b(j)$, $z_3 = b'(k)$, $z_2 = b(i) = b'(i)$. Then $z_0 = \pi(z_1) = x_{\max b(j)}$ and $z_4 = \pi(z_3) = x_{\max b'(k)}$. Observe also that $\delta(z_1, z_2) = d^{\mathbf{U}_Q}(y_j, y_i)$ and that $\delta(z_2, z_3) = d^{\mathbf{U}_Q}(y_i, y_k)$. So:

$$\begin{aligned} \delta(z_0, z_4) &= d^{\mathbf{X}_q}(x_{\max b(j)}, x_{\max b'(k)}) \\ &\leq d^{\mathbf{X}_q}(x_{\max b(j)}, x_{\max b(i)}) + d^{\mathbf{X}_q}(x_{\max b'(i)}, x_{\max b'(k)}) \\ &= d^{\mathbf{X}_q}(x_j, x_i) + d^{\mathbf{X}_q}(x_i, x_k) \\ &= \lceil d^{\mathbf{U}_Q}(y_j, y_i) \rceil_q + \lceil d^{\mathbf{U}_Q}(y_i, y_k) \rceil_q \\ &= \lceil \delta(z_1, z_2) \rceil_q + \lceil \delta(z_2, z_3) \rceil_q \\ &\leq \delta(z_1, z_2) + \frac{1}{q} + \delta(z_2, z_3) + \frac{1}{q} \\ &= \delta(z_1, z_2) + \delta(z_2, z_3) + \frac{2}{q}. \quad \square \end{aligned}$$

3.3. Proof of Lemma 7

Using compactness of \mathbf{K} , find a finite subspace \mathbf{Z} of \mathbf{K} such that $K \subset (Z)_{\varepsilon/2}$.

Claim 3. The space \mathbf{K} embeds in $(\tilde{Z})_{\varepsilon}$ for every subspace \tilde{Z} of \mathbf{U} isometric to \mathbf{Z} .

Proof. This follows from ultrahomogeneity of \mathbf{U} : if \tilde{Z} is a subspace of \mathbf{U} isometric to \mathbf{Z} , let $\phi : Z \rightarrow \tilde{Z}$ be an isometry. By ultrahomogeneity of \mathbf{U} , find $\Phi : U \rightarrow U$ extending ϕ . Then $\Phi(K)$ is isometric to \mathbf{K} and is included in

$$\Phi((Z)_{\varepsilon/2}) = (\Phi(Z))_{\varepsilon/2} = (\tilde{Z})_{\varepsilon/2}. \quad \square$$

Therefore, the space \mathbf{Z} is almost as required except that it may not have rational distances. To arrange that, consider $q \in \mathbb{N}$ large enough so that $1/q < \varepsilon/2$. Recall that for a number α , $\lceil \alpha \rceil_q$ denotes the smallest number $\geq \alpha$ of the form l/q with l integer. The function $\lceil \cdot \rceil_q$ is subadditive and nondecreasing. Hence, the composition $\lceil d^Z \rceil_q = \lceil \cdot \rceil_q \circ d^Z$ is a metric on Z . Let \mathbf{Y} be defined as the metric space $(Z, \lceil d^Z \rceil_q)$. It obviously has rational distances. We are going to show that it is as required. Consider the set $X = Z \times \{0, 1\}$ and define

$$\delta((z, i), (z', i')) = \begin{cases} d^Z(z, z') & \text{if } i = i' = 0, \\ \lceil d^Z(z, z') \rceil_q & \text{if } i = i' = 1, \\ d^Z(z, z') + \varepsilon/2 & \text{if } i \neq i'. \end{cases}$$

In spirit, the structure (X, δ) is obtained by putting a copy of $\mathbf{Y}(= (Z, \lceil d^Z \rceil_q))$ above a copy of \mathbf{Z} such that the distance between any point $(z, 0) \in Z \times \{0\}$ and its counterpart $(z, 1)$ in $Z \times \{1\}$ is $\varepsilon/2$.

Claim 4. The map δ is a metric on X .

Proof. The maps d^Z and $\lceil d^Z \rceil_q$ being metrics on $Z \times \{0\}$ and $Z \times \{1\}$, it suffices to verify that the triangle inequality is satisfied on triangles of the form $\{(x, 0), (y, 0), (z, 1)\}$ and $\{(x, 1), (y, 1), (z, 0)\}$, with $x, y, z \in Z$.

Assume that $x, y, z \in Z$, and consider the triangle $\{(x, 1), (y, 1), (z, 0)\}$. Then

$$\begin{aligned} \delta((x, 1), (z, 0)) &= d^Z(x, z) + \frac{\varepsilon}{2} \\ &\leq d^Z(x, y) + d^Z(y, z) + \frac{\varepsilon}{2} \\ &\leq \lceil d^Z(x, y) \rceil_q + d^Z(y, z) + \frac{\varepsilon}{2} \\ &\leq \delta((x, 1), (y, 1)) + \delta((y, 1), (z, 0)). \end{aligned}$$

Similarly,

$$\delta((y, 1), (z, 0)) \leq \delta((y, 1), (x, 1)) + \delta((x, 1), (z, 0)).$$

And finally,

$$\begin{aligned} \delta((x, 1), (y, 1)) &= \lceil d^Z(x, y) \rceil_q \\ &\leq d^Z(x, y) + \frac{1}{q} \\ &\leq d^Z(x, y) + \frac{\varepsilon}{2} \\ &\leq d^Z(x, z) + d^Z(z, y) + \frac{\varepsilon}{2} \\ &\leq d^Z(x, z) + \frac{\varepsilon}{2} + d^Z(z, y) + \frac{\varepsilon}{2} \\ &\leq \delta((x, 1), (z, 0)) + \delta((z, 0), (y, 1)). \end{aligned}$$

Next, consider the triangle $\{(x, 0), (y, 0), (z, 1)\}$. We have

$$\begin{aligned}\delta((x, 0), (z, 1)) &= d^Z(x, z) + \frac{\varepsilon}{2} \\ &\leq d^Z(x, y) + d^Z(y, z) + \frac{\varepsilon}{2} \\ &\leq \delta((x, 0), (y, 0)) + \delta((y, 0), (z, 1)).\end{aligned}$$

Similarly,

$$\delta((y, 0), (z, 1)) \leq \delta((y, 0), (x, 0)) + \delta((x, 0), (z, 1)).$$

Finally,

$$\begin{aligned}\delta((x, 0), (y, 0)) &= d^Z(x, y) \\ &\leq d^Z(x, z) + d^Z(z, y) \\ &\leq d^Z(x, z) + \frac{\varepsilon}{2} + d^Z(z, y) + \frac{\varepsilon}{2} \\ &\leq \delta((x, 0), (z, 1)) + \delta((z, 1), (y, 0)). \quad \square\end{aligned}$$

Denote the space (X, δ) by \mathbf{X} . Recall that every finite metric space embeds isometrically in \mathbf{U} . Hence, without loss of generality, we may suppose $Y \subset X \subset U$. We claim that \mathbf{Y} is as required. By construction, the space \mathbf{Y} is a finite subspace of \mathbf{U} with distances in \mathbb{N}/q . Observe that $X \subset (Y)_{\varepsilon/2}$. Assume that a subspace $\tilde{\mathbf{Y}}$ of \mathbf{U} is isometric to \mathbf{Y} . By an argument similar to the one used in [Claim 3](#), the space \mathbf{X} embeds in $(\tilde{\mathbf{Y}})_{\varepsilon/2}$. Thus, because \mathbf{Z} embeds in \mathbf{X} , the set $(\tilde{\mathbf{Y}})_{\varepsilon/2}$ contains a copy of \mathbf{Z} , call it $\tilde{\mathbf{Z}}$. By [Claim 3](#), the set $(\tilde{\mathbf{Z}})_{\varepsilon/2}$ contains a copy of \mathbf{K} , call it $\tilde{\mathbf{K}}$. Then

$$\tilde{\mathbf{K}} \subset (\tilde{\mathbf{Z}})_{\varepsilon/2} \subset ((\tilde{\mathbf{Y}})_{\varepsilon/2})_{\varepsilon/2} \subset (\tilde{\mathbf{Y}})_{\varepsilon}.$$

This finishes the proof of [Lemma 7](#).

4. Proof of Theorem 2

In this section, we think of $\mathbf{U}_{\mathbb{Q}}$ as a dense metric subspace of \mathbf{U} . We fix a partition of $\mathbf{U}_{\mathbb{Q}}$ as well as $\varepsilon > 0$, and we assume that there is a finite metric subspace \mathbf{Y} of $\mathbf{U}_{\mathbb{Q}}$ that does not embed in B . Our goal is to show that $\mathbf{U}_{\mathbb{Q}}$ embeds in $(R)_{\varepsilon}$. We start with the following technical lemma:

Lemma 14. *Let \mathbf{V} be a countable subspace of \mathbf{U} with rational distances. Then for every $\varepsilon > 0$ the subspace $\mathbf{U}_{\mathbb{Q}} \cap (\mathbf{V})_{\varepsilon}$ includes a copy of \mathbf{V} .*

Assuming this result for the moment, here is how we prove [Theorem 2](#): let $U_{\mathbb{Q}} = B \cup R$ and $\varepsilon > 0$, and assume that there is a finite metric subspace \mathbf{Y} of $\mathbf{U}_{\mathbb{Q}}$ that does not embed in B . We wish to show that $\mathbf{U}_{\mathbb{Q}}$ embeds in $(R)_{\varepsilon}$. Choose $q \in \mathbb{N}$ large enough so that $2/q \leq \varepsilon$ and all distances appearing in \mathbf{Y} are integer multiples of $1/q$. Working in \mathbf{U} , set

$$\begin{aligned}B^* &= \{x \in U : (\{x\})_{1/2q} \cap U_{\mathbb{Q}} \subset B\}, \\ R^* &= U \setminus B^* = U \cap (R)_{1/2q}.\end{aligned}$$

Consider the space $U_{\mathbb{N}/q}^*$ coming from [Lemma 6](#). The partition $U = B^* \cup R^*$ induces a partition of $U_{\mathbb{N}/q}^*$. Observe that by weak indivisibility of $\mathbf{U}_{\mathbb{N}}$, the space $\mathbf{U}_{\mathbb{N}/q}$ is weakly indivisible as well. We also claim that the space \mathbf{Y} does not embed in $U_{\mathbb{N}/q}^* \cap B^*$. Indeed, otherwise, we could find a copy $\tilde{\mathbf{Y}}$ of \mathbf{Y} in $U_{\mathbb{N}/q}^* \cap B^*$. [Lemma 14](#) applied to $\mathbf{V} = \mathbf{Y}$ would then guarantee that $U_{\mathbb{Q}} \cap (\tilde{\mathbf{Y}})_{1/2q}$ contains a copy of \mathbf{Y} . But by construction, $U_{\mathbb{Q}} \cap (\tilde{\mathbf{Y}})_{1/2q} \subset B$. So \mathbf{Y} would embed in B , a contradiction. Therefore, by weak indivisibility of $\mathbf{U}_{\mathbb{N}/q}$, there is a subspace \mathbf{T} of $U_{\mathbb{N}/q}^*$ isometric to $\mathbf{U}_{\mathbb{N}/q}$ such that $T \subset R^*$. By construction

of $\mathbf{U}_{\mathbb{N}/q}^*$, the set $(T)_{1/q}$ includes an isometric copy of \mathbf{U} , hence an isometric copy $\tilde{\mathbf{U}}$ of $\mathbf{U}_{\mathbb{Q}}$. By Lemma 14 applied to $\mathbf{V} = \tilde{\mathbf{U}}$, the set $U_{\mathbb{Q}} \cap (\tilde{U})_{1/2q}$ contains a copy of $\mathbf{U}_{\mathbb{Q}}$. Observe now that:

$$U_{\mathbb{Q}} \cap (\tilde{U})_{1/2q} \subset ((T)_{1/q})_{1/2q} \subset ((R^*)_{1/q})_{1/2q} \subset (((R)_{1/2q})_{1/q})_{1/2q} \subset (R)_{2/q} \subset (R)_{\varepsilon}.$$

Theorem 2 is proved. The rest of this section is therefore devoted to a proof of Lemma 14.

Proof of Lemma 14. The proof of this lemma closely follows the proof of [6] Proposition 2, and is included here for completeness. We construct the required copy of \mathbf{V} inductively. Consider $\{y_n : n \in \mathbb{N}\}$ an enumeration of \mathbf{V} . For $k \in \mathbb{N}$, set

$$\delta_k = \frac{\varepsilon}{2} \sum_{i=0}^k \frac{1}{2^i}.$$

Set also

$$\eta_k = \frac{\varepsilon}{3} \frac{1}{2^{k+1}}.$$

$\mathbf{U}_{\mathbb{Q}}$ being dense in \mathbf{U} , choose $z_0 \in \mathbf{U}_{\mathbb{Q}}$ such that $d^{\mathbf{U}}(y_0, z_0) < \delta_0$. Assume now that $z_0, \dots, z_n \in \mathbf{U}_{\mathbb{Q}}$ were constructed such that for every $k, l \leq n$

$$\begin{cases} d^{\mathbf{U}}(z_k, z_l) = d^{\mathbf{U}}(y_k, y_l), \\ d^{\mathbf{U}}(z_k, y_k) < \delta_k. \end{cases}$$

Again by denseness of $\mathbf{U}_{\mathbb{Q}}$ in \mathbf{U} , fix $z \in \mathbf{U}_{\mathbb{Q}}$ such that

$$d^{\mathbf{U}}(z, y_{n+1}) < \eta_{n+1}.$$

Then for every $k \leq n$,

$$\begin{aligned} |d^{\mathbf{U}}(z, z_k) - d^{\mathbf{U}}(y_{n+1}, y_k)| &= |d^{\mathbf{U}}(z, z_k) - d^{\mathbf{U}}(z_k, y_{n+1}) + d^{\mathbf{U}}(z_k, y_{n+1}) - d^{\mathbf{U}}(y_{n+1}, y_k)| \\ &\leq d^{\mathbf{U}}(z, y_{n+1}) + d^{\mathbf{U}}(z_k, y_k) \\ &< \eta_{n+1} + \delta_k \\ &< \eta_{n+1} + \delta_n. \end{aligned}$$

It follows that there is $z_{n+1} \in \mathbf{U}_{\mathbb{Q}}$ such that

$$\begin{cases} \forall k \leq n \quad d^{\mathbf{U}}(z_{n+1}, z_k) = d^{\mathbf{U}}(y_{n+1}, y_k) \\ d^{\mathbf{U}}(z_{n+1}, z) < \eta_{n+1} + \delta_n. \end{cases}$$

Indeed, consider the map f defined on $\{z_k : k \leq n\} \cup \{z\}$ by:

$$\begin{cases} \forall k \leq n \quad f(z_k) = d^{\mathbf{U}}(y_{n+1}, y_k), \\ f(z) = \max\{|d^{\mathbf{U}}(z, z_k) - d^{\mathbf{U}}(y_{n+1}, y_k)| : k \leq n\}. \end{cases}$$

Claim 5. f is Katětov.

Proof. The metric space $\{y_k : k \leq n+1\}$ witnesses that f is Katětov over the set $\{z_k : k \leq n\}$ so it suffices to prove that for every $k \leq n$,

$$|f(z) - f(z_k)| \leq d^{\mathbf{U}}(z, z_k) \leq f(z) + f(z_k).$$

Equivalently, for every $k \leq n$,

$$|d^{\mathbf{U}}(z, z_k) - f(z_k)| \leq f(z) \leq d^{\mathbf{U}}(z, z_k) + f(z_k).$$

There is nothing to do for the left-hand side because by definition of f , we have

$$f(z) = \max\{|d^{\mathbf{U}}(z, z_k) - f(z_k)| : k \leq n\}.$$

For right-hand side, what we need to show is that for every $k, l \leq n$,

$$|d^{\mathbf{U}}(z, z_l) - d^{\mathbf{U}}(y_{n+1}, y_l)| \leq d^{\mathbf{U}}(z, z_k) + d^{\mathbf{U}}(y_{n+1}, y_k).$$

Equivalently,

$$\begin{cases} d^{\mathbf{U}}(z, z_l) - d^{\mathbf{U}}(y_{n+1}, y_l) \leq d^{\mathbf{U}}(z, z_k) + d^{\mathbf{U}}(y_{n+1}, y_k), \\ d^{\mathbf{U}}(y_{n+1}, y_l) - d^{\mathbf{U}}(z, z_l) \leq d^{\mathbf{U}}(z, z_k) + d^{\mathbf{U}}(y_{n+1}, y_k). \end{cases}$$

The first inequality is equivalent to

$$d^{\mathbf{U}}(z, z_l) - d^{\mathbf{U}}(z, z_k) \leq d^{\mathbf{U}}(y_{n+1}, y_k) + d^{\mathbf{U}}(y_{n+1}, y_l).$$

But this is satisfied because

$$d^{\mathbf{U}}(z, z_l) - d^{\mathbf{U}}(z, z_k) \leq d^{\mathbf{U}}(z_l, z_k) = d^{\mathbf{U}}(y_k, y_l) \leq d^{\mathbf{U}}(y_k, y_{n+1}) + d^{\mathbf{U}}(y_{n+1}, y_l).$$

Similarly, the second inequality is equivalent to

$$d^{\mathbf{U}}(y_{n+1}, y_l) - d^{\mathbf{U}}(y_{n+1}, y_k) \leq d^{\mathbf{U}}(z, z_k) + d^{\mathbf{U}}(z, z_l).$$

This holds because

$$d^{\mathbf{U}}(y_{n+1}, y_l) - d^{\mathbf{U}}(y_{n+1}, y_k) \leq d^{\mathbf{U}}(y_k, y_l) = d^{\mathbf{U}}(z_k, z_l) \leq d^{\mathbf{U}}(z, z_k) + d^{\mathbf{U}}(z, z_l). \quad \square$$

The map f being Katětov, consider a point $z_{n+1} \in \mathbf{U}_{\mathbb{Q}}$ realizing f over the set $\{z_k : k \leq n\} \cup \{z\}$. Observe then that

$$\begin{aligned} d^{\mathbf{U}}(z_{n+1}, y_{n+1}) &\leq d^{\mathbf{U}}(z_{n+1}, z) + d^{\mathbf{U}}(z, y_{n+1}) \\ &< \eta_{n+1} + \delta_n + \eta_{n+1} \\ &< \delta_{n+1}. \end{aligned}$$

After infinitely many steps, we are left with $\{z_n : n \in \mathbb{N}\} \subset \mathbf{U}_{\mathbb{Q}} \cap (V)_{\varepsilon}$ isometric to \mathbf{V} . \square

5. Age-indivisibility may not imply weak indivisibility

In what follows, the set S is a fixed dense subset of $[0, 2]$. Let \mathcal{E}_S be the class of all finite metric spaces \mathbf{X} with distances in S which embed isometrically into the unit sphere \mathbb{S}^{∞} of ℓ_2 with the property that $\{0_{\ell_2}\} \cup \mathbf{X}$ is affinely independent.

Claim 6. *There is a unique countable ultrahomogeneous metric space \mathbb{S}_S^{∞} whose class of finite metric subspaces is exactly \mathcal{E}_S . Moreover, the metric completion of \mathbb{S}_S^{∞} is \mathbb{S}^{∞} .*

Proof. See [9] or [10]. \square

We show:

Theorem 6. *The space \mathbb{S}_S^{∞} is age-indivisible.*

We also indicate why the space \mathbb{S}_S^{∞} may not be weakly indivisible. The proof of those results are provided in Sections 5.1 and 5.2 respectively.

5.1. The space \mathbb{S}_S^{∞} is age-indivisible

Let \mathbf{Y} be a finite metric subspace of \mathbb{S}_S^{∞} . We need to show:

Claim 7. *There is a finite metric subspace \mathbf{Z} of \mathbb{S}_S^{∞} such that for every partition $Z = B \cup R$, the space \mathbf{Y} embeds in B or R .*

The main ingredient of the proof is the following deep result due to Matoušek and Rödl:

Theorem 7 (Matoušek–Rödl [7]). *Let \mathbf{X} be an affinely independent finite metric subspace of \mathbb{S}^{∞} with circumradius r , and let $\alpha > 0$. Then there is a finite metric subspace \mathbf{Z} of \mathbb{S}^{∞} with circumradius $r + \alpha$ such that for every partition $Z = B \cup R$, the space \mathbf{X} embeds in B or R .*

What we need to prove is that in the case where $\mathbf{X} = \mathbf{Y}$, we may arrange \mathbf{Z} to be a subspace of \mathbb{S}_S^∞ (that is, with distances in S and $\{0_{\ell_2}\} \cup \mathbf{Z}$ affinely independent). We will make use of the following facts along the way:

Theorem 8 (Schoenberg [14]). Let $X = \{x_k : 1 \leq k \leq n\}$ be a finite set and let $\delta : X^2 \rightarrow \mathbb{R}$ satisfying:

- (i) for every $x \in X$, $\delta(x, x) = 0$,
- (ii) for every $x, x' \in X$, $\delta(x, x) = 0$ and $\delta(x', x) = \delta(x, x')$.

Then (X, δ) is isometric to a subset of ℓ_2 iff

$$\max \left\{ \sum_{1 \leq i < j \leq n} \delta(x_i, x_j)^2 \lambda_i \lambda_j : \sum_{k=1}^n \lambda_k^2 = 1 \text{ and } \sum_{k=1}^n \lambda_k = 0 \right\} \leq 0.$$

Moreover, (X, δ) is isometric to an affinely independent subset of ℓ_2 iff the inequality is strict.

Claim 8. Let \mathbf{X} be a finite affinely independent metric subspace of \mathbb{S}^∞ with circumradius r . Then there is $\varepsilon > 0$ such that for every $\delta : X^2 \rightarrow \mathbb{R}$ satisfying

- (i) for every $x, x' \in X$, $\delta(x, x) = 0$ and $\delta(x', x) = \delta(x, x')$,
- (ii) $|\delta^2 - (d^{\mathbf{X}})^2| < \varepsilon^2$,

the space (X, δ) is an affinely independent metric subspace of \mathbb{S}^∞ .

Proof. Direct from [Theorem 8](#) and from the fact that the map $M \mapsto Q_M$ is continuous, where for a matrix $M = (m_{ij})_{1 \leq i, j \leq n}$,

$$Q_M = \max \left\{ \sum_{1 \leq i < j \leq n} m_{ij} \lambda_i \lambda_j : \sum_{k=1}^n \lambda_k^2 = 1 \text{ and } \sum_{k=1}^n \lambda_k = 0 \right\}. \quad \square$$

Claim 9. Let \mathbf{X} be a finite metric subspace of \mathbb{S}^∞ with circumradius r . Let $\varepsilon > 0$. Then $(X, \sqrt{(d^{\mathbf{X}})^2 + \varepsilon^2})$ is Euclidean, affinely independent with circumradius at most $r + \varepsilon$.

Proof. Let V be the affine space spanned by X . Choose $(e_x)_{x \in X}$ a family of pairwise orthogonal unit vectors in V^\perp . For $x \in X$, set $\tilde{x} = x + \varepsilon/\sqrt{2}e_x$. Then the set $\{\tilde{x} : x \in X\}$ is affinely independent and is isometric to $(X, \sqrt{(d^{\mathbf{X}})^2 + \varepsilon^2})$. Its circumradius is at most $r + \varepsilon$ because it is contained in the ball centered at the circumcenter of X and with radius $r + \varepsilon$. \square

Claim 10. Let \mathbf{X} be an affinely independent subspace of \mathbb{S}^∞ . Then $\mathbf{X} \cup \{0_{\ell_2}\}$ is affinely independent iff the circumradius of \mathbf{X} is < 1 .

Proof. Let V be the affine space spanned by X . Then the set $\mathbb{S}^\infty \cap V$ is the circumscribed sphere of X in V . It has radius < 1 iff 0_{ℓ_2} does not belong to V . \square

Proof of Claim 7. First, we show that there is an affinely independent finite metric subspace \mathbf{Z}_0 of \mathbb{S}^∞ with circumradius < 1 such that for every partition $Z_0 = B \cup R$, \mathbf{Y} embeds in B or R :

Let r denote the circumradius of \mathbf{Y} . Because \mathbf{Y} is a subspace of \mathbb{S}_S^∞ , the space $\mathbf{Y} \cup \{0_{\ell_2}\}$ is affinely independent and by [Claim 10](#), we have $r < 1$. By [Claim 8](#), fix $\varepsilon > 0$ such that $r + 2\varepsilon < 1$ and such that for every map $\delta : Y^2 \rightarrow \mathbb{R}$ satisfying

- (i) for every $y, y' \in Y$, $\delta(y, y) = 0$ and $\delta(y', y) = \delta(y, y')$,
- (ii) $|\delta^2 - (d^{\mathbf{Y}})^2| < \varepsilon^2$,

the space (Y, δ) is still Euclidean and affinely independent. Fix $\alpha > 0$ such that $\alpha < \varepsilon$. By choice

of α , the space $(Y, \sqrt{(d^Y)^2 - \alpha^2})$ is still Euclidean and affinely independent. It should be clear that its circumradius is at most r . Apply [Theorem 7](#) to produce a finite metric subspace \mathbf{T} of \mathbb{S}^∞ with circumradius $r + \alpha$ such that for every partition $T = B \cup R$, the space $(Y, \sqrt{(d^Y)^2 - \alpha^2})$ embeds in B or R . Set $\mathbf{Z}_0 = (T, \sqrt{(d^T)^2 + \alpha^2})$. We claim that \mathbf{Z}_0 is as required.

Indeed, by [Claim 9](#), \mathbf{Z}_0 is Euclidean, affinely independent, and its circumradius is at most $(r + \alpha) + \alpha < r + 2\varepsilon < 1$. Next, if $\mathbf{Z}_0 = B \cup R$, this partition induces a partition $T = B \cup R$. By construction of \mathbf{T} , there is a subspace $\tilde{\mathbf{Y}}$ of \mathbf{T} isometric to $(Y, \sqrt{(d^Y)^2 - \alpha^2})$ contained in B or R . Note that in \mathbf{Z}_0 , the metric subspace supported by $\tilde{\mathbf{Y}}$ is isometric to

$$\left(Y, \sqrt{\left(\sqrt{(d^Y)^2 - \alpha^2} \right)^2 + \alpha^2} \right) = \left(Y, \sqrt{(d^Y)^2 - \alpha^2 + \alpha^2} \right) = (Y, d^Y) = \mathbf{Y}.$$

Consider the space \mathbf{Z}_0 we just constructed. Using [Claim 8](#) as well as the denseness of S , we may modify slightly all the distances in \mathbf{Z}_0 that are not in S and turn \mathbf{Z}_0 into an affinely independent subspace \mathbf{Z} of \mathbb{S}^∞ with distances in S and circumradius < 1 . By [Claim 10](#), the space $\{0_{\ell_2}\} \cup \mathbf{Z}$ is affinely independent. Therefore, \mathbf{Z} embeds in \mathbb{S}_S^∞ . Finally, note that since all the distances of \mathbf{Z}_0 that were already in S did not get changed, the copies of \mathbf{Y} in \mathbf{Z}_0 remain unaltered when passing to \mathbf{Z} . It follows that for every partition $Z = B \cup R$, the space \mathbf{Y} embeds in B or R . \square

5.2. The space \mathbb{S}_S^∞ may not be weakly indivisible

The starting point of this section is the following theorem:

Theorem 9 (Odell–Schlumprecht [12]). *There is a partition $\mathbb{S}^\infty = B \cup R$ and $\varepsilon > 0$ such that*

- (i) *For every linear subspace V of ℓ_2 with $\dim V = \infty$, $\mathbb{S}^\infty \cap V \not\subset (B)_\varepsilon$.*
- (ii) *For every linear subspace V of ℓ_2 with $\dim V = \infty$, $\mathbb{S}^\infty \cap V \not\subset (R)_\varepsilon$.*

In response to an inquiry of the authors, Thomas Schlumprecht [13] indicated that the method that was used to prove Theorem 9 in [12] (where the statement is proved first in another Banach space known as the Schlumprecht space, and then transferred to ℓ_2), can be adapted to show that $\dim V = \infty$ may be replaced by $\dim V = 2$ in (i). However, he indicated recently that some obstruction had appeared. Nevertheless, we would like to present here how the aforementioned strengthening of [Theorem 9](#) implies that \mathbb{S}_S^∞ is not weakly indivisible.

Theorem 10. *Assume that there is a partition $\mathbb{S}^\infty = B \cup R$ and $\varepsilon > 0$ such that*

- (i) *for every linear subspace V of ℓ_2 with $\dim V = 2$, $\mathbb{S}^\infty \cap V \not\subset (B)_\varepsilon$,*
- (ii) *for every linear subspace V of ℓ_2 with $\dim V = \infty$, $\mathbb{S}^\infty \cap V \not\subset (R)_\varepsilon$.*

Then \mathbb{S}_S^∞ is not weakly indivisible.

Consider the partition of \mathbb{S}^∞ provided by [Theorem 10](#). It should be clear that it induces a partition of \mathbb{S}_S^∞ .

Claim 11. $\mathbb{S}_S^\infty = B \cup R$ witnesses that \mathbb{S}_S^∞ is not weakly indivisible.

The proof makes use of the following fact, which we prove for completeness:

Claim 12. *Let $Y \subset \mathbb{S}^\infty$ be isometric to \mathbb{S}^∞ . Then there is a closed linear subspace V of ℓ_2 with $\dim V = \infty$ such that $Y = V \cap \mathbb{S}^\infty$.*

Proof. Consider V the closed linear span of Y in ℓ_2 . Consider also the set $W = \{\lambda y : \lambda \in \mathbb{R}, y \in Y\}$. We will be done if we show $V = W$. Clearly, $W \subset V$. For the reverse inclusion, observe that because Y is closed (it is isometric to a complete metric space), the set W is closed. Therefore, it is enough to

show that all the finite linear combinations of elements of V that have norm 1 are in Y , i.e. for every $y_1, \dots, y_n \in Y$ and $\lambda_1, \dots, \lambda_n \in \mathbb{R}$ such that $\sum_{i=1}^n \lambda_i y_i \neq 0_{\ell_2}$,

$$\frac{\sum_{i=1}^n \lambda_i y_i}{\left\| \sum_{i=1}^n \lambda_i y_i \right\|} \in Y.$$

We proceed by induction on n . For $n = 2$, we first consider the case $\lambda_1 = \lambda_2 = 1$. Note that y_1 and y_2 cannot be antipodal (otherwise $y_1 + y_2 = 0_{\ell_2}$), and that $\frac{y_1 + y_2}{\|y_1 + y_2\|}$ can be characterized metrically in terms of y_1 and y_2 . For example, it is the unique geodesic middle point of y_1 and y_2 in the intrinsic metric on \mathbb{S}^∞ . Since the intrinsic metric can be defined in terms of the usual Hilbertian metric on \mathbb{S}^∞ , this point must belong to Y . By a usual middle-point-type argument, it follows that the entire geodesic segment between y_1 and y_2 is contained in Y . Using then that Y is closed under antipodality (because Y being isometric to \mathbb{S}^∞ any $y \in Y$ must have a point at distance 2), as well as a middle-point-type argument again, the entire great circle through y_1 and y_2 is contained in Y . That finishes the case $n = 2$. Assume that the property is proved up to $n \geq 2$. Fix $y_1, \dots, y_{n+1} \in Y$ and $\lambda_1, \dots, \lambda_{n+1} \in \mathbb{R}$. Then writing

$$z = \frac{\sum_{i=1}^n \lambda_i y_i}{\left\| \sum_{i=1}^n \lambda_i y_i \right\|},$$

the vector

$$\frac{\sum_{i=1}^{n+1} \lambda_i y_i}{\left\| \sum_{i=1}^{n+1} \lambda_i y_i \right\|}$$

is a linear combination of z and y_{n+1} with norm 1. Therefore, it is of the form

$$\frac{\alpha z + \beta y_{n+1}}{\|\alpha z + \beta y_{n+1}\|}.$$

By induction hypothesis, z is in Y . So again by induction hypothesis (case $n = 2$),

$$\frac{\alpha z + \beta y_{n+1}}{\|\alpha z + \beta y_{n+1}\|} \in Y.$$

Therefore,

$$\frac{\sum_{i=1}^{n+1} \lambda_i y_i}{\left\| \sum_{i=1}^{n+1} \lambda_i y_i \right\|} \in Y. \quad \square$$

Proof of Claim 11. Let W be a linear subspace of ℓ_2 with $\dim W = 2$. By compactness of $\mathbb{S}^\infty \cap W$ and denseness of \mathbb{S}_S^∞ in \mathbb{S}^∞ , there is $X \subset \mathbb{S}_S^\infty$ finite such that $\mathbb{S}^\infty \cap W \subset (X)_\varepsilon$. Let \mathbf{X} denote the metric subspace of \mathbb{S}_S^∞ supported by the set X . Then \mathbf{X} does not embed in B because otherwise, there would be a linear subspace V of ℓ_2 with $\dim V = 2$ such that $\mathbb{S}^\infty \cap V \subset (B)_\varepsilon$, violating (i) of [Theorem 10](#). On the other hand, \mathbb{S}_S^∞ cannot embed in R : let $Y \subset \mathbb{S}_S^\infty$ be isometric to \mathbb{S}_S^∞ . Then in \mathbb{S}^∞ , the closure \bar{Y} of Y is isometric to \mathbb{S}^∞ . By [Claim 11](#), there is a closed linear subspace V of ℓ_2 with $\dim V = \infty$ such that $\bar{Y} = V \cap \mathbb{S}^\infty$. By (ii) of [Theorem 10](#), $\bar{Y} \not\subset (R)_\varepsilon$. Therefore $\bar{Y} \not\subset R$. \square

Acknowledgements

We would like to thank Thomas Schlumprecht for helpful discussions concerning the potential negation of weak indivisibility of $\mathbb{S}_{\mathbb{Q}}^{\infty}$. We would also like to express our gratitude to three anonymous referees, thanks to whom the quality of this paper considerably improved. L. Nguyen Van Thé would like to acknowledge the support of the Department of Mathematics & Statistics Postdoctoral Program at the University of Calgary, as well as the support of the Swiss National Fund via the postdoctoral fellowship # 20-118014-1. N.W. Sauer was supported by NSERC of Canada Grant # 691325.

References

- [1] C. Delhommé, C. Laflamme, M. Pouzet, N. Sauer, Divisibility of countable metric spaces, *European J. Combin.* 28 (6) (2007) 1746–1769.
- [2] R. Fraïssé, *Theory of Relations*, in: *Studies in Logic and the Foundations of Mathematics*, vol. 145, North-Holland Publishing Co., Amsterdam, 2000.
- [3] A.S. Kechris, V. Pestov, S. Todorcevic, Fraïssé limits, Ramsey theory, and topological dynamics of automorphism groups, *Geom. Funct. Anal.* 15 (2005) 106–189.
- [4] C. Laflamme, L. Nguyen Van Thé, M. Pouzet, N.W. Sauer, Partitions and indivisibility properties of countable infinite dimensional vector spaces, *J. Combin. Theory Ser. A* (2010) doi:10.1016/j.jcta.2009.10.007.
- [5] A. Leiderman, V. Pestov, M. Rubin, S. Solecki, V.V. Uspenskij (Eds.), *Workshop on the Urysohn space*, Ben-Gurion University of the Negev, Beer Sheva, May 21–24, 2006, *Topology Appl.* 155 (14) (2008) (special issue).
- [6] J. Lopez-Abad, L. Nguyen Van Thé, The oscillation stability problem for the Urysohn sphere: A combinatorial approach, *Topology Appl.* 155 (14) (2008) 1516–1530.
- [7] J. Matoušek, V. Rödl, On Ramsey sets in spheres, *J. Combin. Theory Ser. A* 70 (1995) 30–44.
- [8] J. Nešetřil, Metric spaces are Ramsey, *European J. Combin.* 28 (1) (2007) 457–468.
- [9] L. Nguyen Van Thé, *Théorie de Ramsey structurale des espaces métriques et dynamique topologique des groupes d'isométries*, Ph.D. Thesis, Université Paris 7, 2006, available in English at: <http://tel.archives-ouvertes.fr/tel-00139239/fr/>.
- [10] L. Nguyen Van Thé, Structural Ramsey theory of metric spaces and topological dynamics of isometry groups, *Mem. Amer. Math. Soc.* (in press) preprint available at: [arXiv:0804.1593](https://arxiv.org/abs/0804.1593).
- [11] L. Nguyen Van Thé, N.W. Sauer, The Urysohn sphere is oscillation stable, *GAFA, Geom. Funct. Anal.* 19 (2009) 536–557.
- [12] E. Odell, T. Schlumprecht, The distortion problem, *Acta Math.* 173 (1994) 259–281.
- [13] T. Schlumprecht, Personal communication.
- [14] I.J. Schoenberg, Metric spaces and positive definite functions, *Trans. Amer. Math. Soc.* 44 (3) (1938) 522–536.